

LONGEST SUBSEQUENCE FOR CERTAIN REPEATED UP/DOWN PATTERNS IN RANDOM PERMUTATIONS AVOIDING A PATTERN OF LENGTH THREE

ROSS G. PINSKY

ABSTRACT. Let S_n denote the set of permutations of $[n]$ and let $\sigma = \sigma_1 \cdots \sigma_n \in S_n$. For any subsequence $\{\sigma_{i_j}\}_{j=1}^k$ of $\{\sigma_i\}_{i=1}^n$ of length $k \geq 2$, construct the “up/down” sequence $V_1 \cdots V_{k-1}$ defined by

$$V_j = \begin{cases} U, & \text{if } \sigma_{i_{j+1}} - \sigma_{i_j} > 0; \\ D, & \text{if } \sigma_{i_{j+1}} - \sigma_{i_j} < 0, \end{cases}$$

where U refers to “up”, D to “down” and V to “vertical”. Consider now a fixed up/down pattern: $V_1 \cdots V_l$, where $l \in \mathbb{N}$ and $V_j \in \{U, D\}$, $j \in [l]$. Given a permutation $\sigma \in S_n$, consider the length of the longest subsequence of σ that repeats this pattern. Incomplete patterns are not counted, so the length is necessarily either 0 or of the form $kl + 1$, where $k \in \mathbb{N}$. For example, consider $l = 3$ and $V_1 V_2 V_3 = UUD$. Then for the permutation $342617985 \in S_9$, the length of the longest subsequence that repeats the pattern UUD is 7; it is obtained by three different subsequences, namely 3461798, 3461795 and 3461785.

The above framework includes one much studied case as well as another case that has been studied to some degree. The pattern U is the celebrated case of the longest increasing subsequence. The pattern UD (or DU) is the case of the longest alternating subsequence. These have been studied both under the uniform distribution on S_n as well as under the uniform distribution on those permutations in S_n which avoid a particular pattern of length three.

In this paper, we consider the patterns UUD and $UUUD$ under the uniform distribution on those permutations in S_n that avoid the pattern 132. We prove that the expected value of the longest increasing subsequence following the pattern UUD is asymptotic to $\frac{3}{7}n$ and the expected value of the longest increasing subsequence following the pattern $UUUD$ is asymptotic to $\frac{4}{11}n$. (For UD (alternating subsequences) it is known to be $\frac{1}{2}n$.) This leads directly to appropriate corresponding results for permutations avoiding any particular pattern of length three.

1. INTRODUCTION AND STATEMENT OF RESULTS

Let S_n denote the set of permutations of $[n] = \{1, \dots, n\}$ and let $\sigma = \sigma_1 \cdots \sigma_n \in S_n$. For any subsequence $\{\sigma_{i_j}\}_{j=1}^k$ of $\{\sigma_i\}_{i=1}^n$ of length $k \geq 2$, construct the “up/down” sequence $V_1 \cdots V_{k-1}$ defined by

$$V_j = \begin{cases} U, & \text{if } \sigma_{i_{j+1}} - \sigma_{i_j} > 0; \\ D, & \text{if } \sigma_{i_{j+1}} - \sigma_{i_j} < 0, \end{cases}$$

where U refers to “up”, D refers to “down” and V refers to “vertical”.

Consider now a fixed up/down pattern: $V_1 \cdots V_l$, where $l \in \mathbb{N}$ and $V_j \in \{U, D\}$, $j \in [l]$. Given a permutation $\sigma \in S_n$, consider the length of the longest subsequence of σ that repeats this pattern. Incomplete patterns are not counted, so the length is necessarily either 0 or of the form $kl + 1$, where $k \in \mathbb{N}$. For example, consider $l = 3$ and $V_1 V_2 V_3 = UUD$. Then for the permutation $342617985 \in S_9$, the length of the longest subsequence that repeats the pattern UUD is 7; it is obtained by three different subsequences, namely 3461798, 3461795 and 3461785. On the other hand, for the permutation 319652478, the length of the longest subsequence that repeats the pattern UUD is 0 because this pattern does not appear at all.

The above framework includes one very celebrated and much studied case as well as another case that has been studied to some degree. The pattern U is the case of the longest increasing subsequence. This celebrated case was studied by Logan and Shepp [7] and Vershik and Kerov [12]. Their work showed that the expected value of the length of the longest increasing subsequence in a uniformly random permutation from S_n behaves asymptotically as $2\sqrt{n}$. More precise information on the behavior of this random variable was obtained later in the seminal paper of Baik, Deift and Johansson [2]; for more on the longest increasing subsequence and many additional references, see the book by Romik [9].

The pattern UD (or DU) is the case of the longest alternating subsequence (in the first case starting with increasing and ending with decreasing, and in the second case vice versa). Stanley [11] investigated alternating sequences and showed that the expected value of the longest alternating subsequence in a uniformly random permutation from S_n behaves asymptotically as $\frac{2}{3}n$. (Of course this asymptotic behavior is independent of how we define the initial or terminal direction in the sequence.) See also further results by Widom [13]. The analysis in the alternating case is simpler than in the increasing case

because, as Stanley noted, there is always a longest alternating subsequence (either beginning with down or ending with up) of $\sigma \in S_n$ which contains the number n . Thus, a longest subsequence can be broken down into smaller pieces that are concatenated.

In [11], Stanley also posed the question of whether it is true that for any pattern of ups and downs as we have defined above, there exist constants μ and c such that the expected value of the length of the longest subsequence repeating this pattern in a uniformly random permutation from S_n behaves asymptotically as μn^c . The recent paper [1] answered this question in the affirmative, and showed in particular that for every pattern except for the pattern U corresponding to the longest increasing subsequence, one has $c = 1$. Thus, for every pattern except for U , the asymptotic behavior of the expected value of the length of the longest subsequence repeating that pattern is μn , for some $\mu \in (0, 1)$. The authors of [1] did not explicitly calculate the value of μ ; however, they constructed a dynamical system that can be used to approximate μ . They also proved a central limit theorem for the length of the longest subsequence repeating any particular pattern, except for the pattern U .

In this paper, instead of considering a permutation that is uniformly random from S_n , we consider a permutation that is uniformly random from those permutations in S_n which avoid a certain pattern of length three. Our aim is to calculate explicitly the asymptotic behavior of the expected value of the length of the longest subsequence repeating certain patterns of ups and downs in such a random permutation. Before continuing to explain this, we recall the definition of pattern-avoiding for permutations. If $\sigma = \sigma_1 \cdots \sigma_n \in S_n$ and $\tau = \tau_1 \cdots \tau_m \in S_m$, where $2 \leq m < n$, then we say that σ contains τ as a pattern if there exists a subsequence $1 \leq i_1 < i_2 < \cdots < i_m \leq n$ such that for all $1 \leq j, k \leq m$, the inequality $\sigma_{i_j} < \sigma_{i_k}$ holds if and only if the inequality $\tau_j < \tau_k$ holds. If σ does not contain τ , then we say that σ avoids τ . We consider here permutations on S_n that avoid a pattern $\tau \in S_3$. Denote by $S_n(\tau)$ the set of permutation in S_n that avoid τ . It is well-known that $|S_n(\tau)| = C_n$, for all six permutations $\tau \in S_3$, where $C_n = \frac{1}{n+1} \binom{2n}{n}$, $n \in \mathbb{N}$, is the n th Catalan number [3]. Let $P_n^{\text{av}(\tau)}$ denote the uniform probability measure on $S_n(\tau)$ and let $E_n^{\text{av}(\tau)}$ denote the corresponding expectation.

As already noted, the pattern U corresponds to increasing subsequences. In [4], the asymptotic behavior of the expectation of the longest increasing

subsequence L_n of a random permutation under the distribution $P_n^{\text{av}(\tau)}$ was obtained for all six permutations $\tau \in S_3$. Of course, the case $\tau = 123$ is trivial. The expectation is on the order n only for $\tau \in \{231, 312, 321\}$. The asymptotic behavior of the variance $v_n(\tau)$ was also investigated, and the limiting distribution of $\frac{L_n - E_n^{\text{av}(\tau)} L_n}{v_n(\tau)}$ was calculated, the limit being Gaussian only for $\tau \in \{231, 312\}$. Large deviations were considered in [8].

As already noted, the pattern UD (or DU) corresponds to alternating subsequences. In [6], the asymptotic behavior of the expectation of the longest alternating sequence of a random permutation under the distribution $P_n^{\text{av}(\tau)}$ was shown to be $\frac{n}{2}$ for all six choices of $\tau \in S_n$. The asymptotic variance was also obtained as well as a central limit theorem. Large deviations were considered in [8].

In this paper, for the patterns UUD and $UUUD$, we will calculate the asymptotic behavior of the expectation of the longest subsequence repeating that pattern in a uniformly random permutation avoiding the pattern 132. The proof in the case of UUD involves analyzing the asymptotic behavior of the coefficients of either of two generating functions that satisfy a system of two linear equations. The calculations are somewhat involved. The proof in the case $UUUD$ involves analyzing the asymptotic behavior of the coefficients of any one of three generating functions that satisfy a system of three linear equations. Here the calculations are quite involved. Using the same method, we could also obtain the asymptotic behavior of the variance, but we have decided not to pursue this, as those calculations would be even more involved. In principle, our technique can be continued for the pattern $U^l D$, for any $l \in \mathbb{N}$, where U^l indicates l consecutive U 's. However, this involves solving a system of l linear equations for l generating functions, solving explicitly for one of them, and then analyzing its coefficients. It also involves the solving of an auxiliary set of equations to calculate the probability that $\sigma \in S_n(132)$ does not have an increasing subsequence of length j , for $j = 1, \dots, l$ (an extension of Lemma 1 in Section 3). The reason the cases $U^l D$ are in principle tractable for 132-avoiding permutations is that a variant of Stanley's observation holds in these cases; namely, that for a 132-avoiding permutation, there is always either a longest subsequence repeating the pattern $U^l D$ that contains the number n , or else, every such longest subsequence starts after the appearance of the number n in the permutation.

Using the reversal, complementation, and reversal-complementation operations for permutations, the results we obtain for permutations avoiding the pattern 132 can be translated into similar results for permutations avoiding any one of the patterns 213, 231, 312. Using a well-known bijection between permutations avoiding the pattern 132 and permutations avoiding the pattern 123, along with reversal, the results we obtain can be translated into similar results for permutations avoiding either of the patterns 123, 321.

We now state two theorems for 132-avoiding permutations, one for the pattern UUD and one for the pattern $UUUD$, and then state a corollary of these two theorems that contains similar results for τ -avoiding permutations for the other five $\tau \in S_3$.

Theorem 1. *Let $L_n^{U^2D}(\sigma)$ denote the length of the longest subsequence of the repeated pattern UUD in $\sigma \in S_n(132)$. (So $L_n^{U^2D}(\sigma)$ is either equal to 0 or to $3k + 1$ for some $k \in \mathbb{N}$.) Then*

$$(1.1) \quad E_n^{av(132)} L_n^{U^2D} \sim \frac{3}{7}n.$$

Theorem 2. *Let $L_n^{U^3D}(\sigma)$ denote the length of the longest subsequence of the repeated pattern $UUUD$ in $\sigma \in S_n(132)$. (So $L_n^{U^3D}(\sigma)$ is either equal to 0 or to $4k + 1$ for some $k \in \mathbb{N}$.) Then*

$$(1.2) \quad E_n^{av(132)} L_n^{U^3D} \sim \frac{4}{11}n.$$

Remark. Recall that it was noted above that for the repeated pattern UD (which corresponds to alternating subsequences), the corresponding asymptotic behavior is $\frac{1}{2}n$.

Corollary 1. *i. Let $L_n^{V_1V_2V_3}(\sigma)$ denote the length of the longest subsequence of the repeated pattern $V_1V_2V_3$ in $\sigma \in S_n$, where $V_i \in \{U, D\}$, $i = 1, 2, 3$. Then*

$$(1.3) \quad E_n^{av(\tau)} L_n^{V_1V_2V_3} \sim \frac{3}{7}n,$$

for the following five pairs of $V_1V_2V_3$ and τ : UDD and 231 ; DDU and 312 ; DUU and 213 ; UDD and 123 ; UUD and 321 .

ii. Let $L_n^{V_1V_2V_3V_4}(\sigma)$ denote the length of the longest subsequence of the repeated pattern $V_1V_2V_3V_4$ in $\sigma \in S_n$, where $V_i \in \{U, D\}$, $i = 1, 2, 3, 4$. Then

$$(1.4) \quad E_n^{av(\tau)} L_n^{V_1V_2V_3V_4} \sim \frac{4}{11}n,$$

for the following five pairs of $V_1V_2V_3$ and τ : $UDDD$ and 231 ; $DDDU$ and 312 ; $DUUU$ and 213 ; $UDDD$ and 123 ; $UUUD$ and 321 .

Proof of Corollary. Recall that the *reverse* of a permutation $\sigma = \sigma_1 \cdots \sigma_n$ is the permutation $\sigma^{\text{rev}} := \sigma_n \cdots \sigma_1$, and the *complement* of σ is the permutation σ^{com} satisfying $\sigma_i^{\text{com}} = n + 1 - \sigma_i$, $i = 1, \dots, n$. Let $\sigma^{\text{rev-com}}$ denote the permutation obtained by applying reversal and then complementation to σ (or equivalently, vice versa). Since $132^{\text{rev}} = 231$, $132^{\text{comp}} = 312$ and $132^{\text{rev-com}} = 213$, it follows that the three operations, reversal, complementation and reversal-complementation, are bijections from $S_n(132)$ to $S_n(\tau)$, with $\tau = 231$ in the case of reversal, $\tau = 312$ in the case of complementation and $\tau = 213$ in the case of reversal-complementation. From these facts and Theorems 1 and 2, the corollary follows immediately for $\tau \in \{231, 312, 213\}$.

There is a well-known explicit bijection between $S_n(132)$ and $S_n(123)$ [10, 3]. Recall that an entry $j \in [n]$ of a permutation $\sigma \in S_n$ is called a *left-to-right minimum* if $\sigma_j = \min\{\sigma_i : 1 \leq i \leq j\}$. For a permutation $\sigma \in S_n(132)$, let $\{i_j\}_{j=1}^k$ denote its left-to-right minima. Then necessarily the entries of σ that appear from left to right between σ_{i_j} and $\sigma_{i_{j+1}}$ (or after σ_{i_k} up through the final term in the permutation) are increasing, with each entry being the smallest number remaining that is larger than its predecessor. (In particular, the left most such entry is the smallest remaining number larger than σ_{i_j} .) The bijection between $S_n(132)$ and $S_n(123)$ preserves the set of left-to-right minima, and then rearranges all of the other entries in descending order from left to right. Note that the values of the permutation at the left-to-right minima form a decreasing sequence, and these other rearranged entries also form a decreasing sequence; thus the permutation obtained is the union of two decreasing sequences, which is equivalent to its being 123-avoiding. One can check easily that for the pattern UUD (or $UUUD$), there is either no copy or one copy of the pattern between two consecutive left-to-right minima in the permutation $\sigma \in S_n(132)$, and that the same number of copies of UDD (or $UDDD$) appear between those two consecutive left-to-right minima in the 123-avoiding permutation obtained from σ via the above described bijection. This proves the corollary for $\tau = 123$. Applying reversal to 123 proves the corollary for $\tau = 321$. \square

We prove Theorem 1 in Section 2. We derive a system of two linear equations for two generating functions, and then solve explicitly for one of them. These generating functions are connected to the expected number of complete patterns UUD in a maximal subsequence. The leading order asymptotic behavior of the coefficients of either of these generating functions

is equal to the leading order asymptotic behavior of $\frac{1}{3}C_n E_n^{\text{av}(132)} L_n^{U^2D}$. Performing an asymptotic analysis of the coefficients of this generating function yields the proof of the theorem.

The proof of Theorem 2 is much longer. In Section 3 we derive a system of three linear equations for three generating functions, and then solve explicitly for one of them. The explicit expression for this generating function is quite involved. These generating functions are connected to the expected number of complete patterns $UUUD$ in a maximal subsequence. The leading order asymptotic behavior of the coefficients of any of these three generating functions is equal to the leading order asymptotic behavior of $\frac{1}{4}C_n E_n^{\text{av}(132)} L_n^{U^3D}$. In Section 4 we perform a lot of algebra in order to obtain the generating function in a more manageable form. Then we perform an asymptotic analysis of the coefficients of this generating function to yield the proof of the theorem.

2. PROOF OF THEOREM 1

For $\sigma \in S_n$ and $n \in \mathbb{N}$, define $B_n^{U^2D}(\sigma)$ to be the number of complete sets of UUD in a longest subsequence in σ of the repeated pattern UUD . Thus,

$$(2.1) \quad B_n^{U^2D}(\sigma) = \begin{cases} \frac{1}{3} \left(L_n^{U^2D}(\sigma) - 1 \right), & \text{if } L_n^{U^2D}(\sigma) \neq 0; \\ 0, & \text{if } L_n^{U^2D}(\sigma) = 0. \end{cases}$$

Also, for convenience, we define $B_0^{U^2D} \equiv 0$.

For $\sigma \in S_n$ and $n \in \mathbb{N}$, define $A_n^{U^2D}(\sigma) = 0$, if $\sigma = n \cdots 21$; otherwise, find a longest subsequence $\{\sigma_{i_j}\}_{j=1}^{3k+2}$, $k \in \mathbb{Z}^+$, of σ for which the up/down pattern is $UUD \cdots UUDU$, and define $A_n^{U^2D}(\sigma) = k + 1$. For convenience, we define $A_0^{U^2D} \equiv 0$.

In the sequel, for any $j \in \mathbb{N}$, $B_j^{U^2D}$ and $A_j^{U^2D}$ will always be considered as random variables on the probability space $(S_j(132), P_j^{\text{av}(132)})$. Define

$$(2.2) \quad b_n = E_n^{\text{av}(132)} B_n^{U^2D}; \quad a_n = E_n^{\text{av}(132)} A_n^{U^2D},$$

where we have suppressed the notation U^2D . Define the generating functions for $\{C_n b_n\}_{n=0}^{\infty}$ and $\{C_n a_n\}_{n=0}^{\infty}$ by

$$(2.3) \quad \begin{aligned} \mathcal{B}^{U^2D}(t) &= \sum_{n=0}^{\infty} C_n b_n t^n; \\ \mathcal{A}^{U^2D}(t) &= \sum_{n=0}^{\infty} C_n a_n t^n. \end{aligned}$$

Also let $C(t) = \sum_{n=0}^{\infty} C_n t^n$ denote the generating function of the Catalan numbers, where we define $C_0 = 1$. As is well-known,

$$(2.4) \quad C(t) = \frac{1 - \sqrt{1 - 4t}}{2t}.$$

The following definition will be useful. Let $a_1 < a_2 < \dots < a_m$ be real numbers and let $\rho = \rho_1 \dots \rho_m$ be a permutation of these numbers. We define $\text{red}(\rho) \in S_m$, the reduction of ρ , to be the permutation in S_m that has the same pattern as ρ . That is, $\text{red}(\rho) = \sigma$ if σ satisfies $\sigma_i < \sigma_j$ whenever $\rho_i < \rho_j$, $i, j \in [m]$. Note that the up/down pattern that one can associate with $\rho = \rho_1 \dots \rho_m$ is the same as the up/down pattern associated with $\text{red}(\rho)$. Every permutation $\sigma \in S_n(132)$ has the property that if $\sigma_j = n$, then the numbers $\{n - j + 1, \dots, n - 1\}$ appear in the first $j - 1$ positions in σ and the numbers $\{1, \dots, n - j\}$ appear in the last $n - j$ positions in σ . From this fact, along with the fact that $|S_n(132)| = C_n$, it follows that

$$(2.5) \quad P_n^{\text{av}(132)}(\sigma_j = n) = \frac{C_{j-1}C_{n-j}}{C_n}, \text{ for } j \in [n].$$

It also follows that under the conditioned measure $P_n^{\text{av}(132)}|_{\{\sigma_j = n\}}$, the permutation $\text{red}(\sigma_1 \dots \sigma_{j-1}) \in S_{j-1}$ has the distribution $P_{j-1}^{\text{av}(132)}$, the permutation $\sigma_{j+1} \dots \sigma_n \in S_{n-j}$ has the distribution $P_{n-j}^{\text{av}(132)}$, and these two permutations are independent.

We now derive a system of two linear equations for $\mathcal{B}^{U^2D}(t)$ and $\mathcal{A}^{U^2D}(t)$, and then solve for one of them explicitly. From the definitions of $B_n^{U^2D}$ and $A_n^{U^2D}$, we have

$$B_n^{U^2D} \equiv 0, \quad 0 \leq n \leq 3; \quad A_n^{U^2D} \equiv 0, \quad 0 \leq n \leq 1.$$

Thus,

$$(2.6) \quad \begin{aligned} b_n &= 0, \quad 0 \leq n \leq 3; \\ a_n &= 0, \quad 0 \leq n \leq 1. \end{aligned}$$

The following proposition is the key to obtaining a pair of linear equations for the generating functions $\mathcal{B}^{U^2D}(t)$ and $\mathcal{A}^{U^2D}(t)$.

Proposition 1. *i.*

$$(2.7) \quad \begin{aligned} B_n^{U^2D}|\{\sigma_j = n\} &\stackrel{\text{dist}}{=} A_{j-1}^{U^2D} + B_{n-j}^{U^2D}, \quad j \in [n-1], \quad n \geq 2; \\ B_n^{U^2D}|\{\sigma_n = n\} &\stackrel{\text{dist}}{=} B_{n-1}^{U^2D}, \quad n \geq 2, \end{aligned}$$

where on the right hand side of (2.7), $A_{j-1}^{U^2D}$ and $B_{n-j}^{U^2D}$ are understood to be independent.

ii.

(2.8)

$$\begin{aligned} A_n^{U^2D}|\{\sigma_1 = n\} &\stackrel{\text{dist}}{=} A_{n-1}^{U^2D}, \quad n \geq 2; \\ A_n^{U^2D}|\{\sigma_j = n\} &\stackrel{\text{dist}}{=} \left(A_{j-1}^{U^2D} + A_{n-j}^{U^2D}\right) 1_{\{A_{n-j}^{U^2D} \neq 0\}} + \left(B_{j-1}^{U^2D} + 1\right) 1_{\{A_{n-j}^{U^2D} = 0\}}, \\ j &\in \{2, \dots, n\}, \quad n \geq 2, \end{aligned}$$

where on the right hand side of (2.8), $A_{j-1}^{U^2D}$ and $A_{n-j}^{U^2D}$ are understood to be independent and $B_{j-1}^{U^2D}$ and $A_{n-j}^{U^2D}$ are understood to be independent.

Proof. The first line of (2.7) follows from the equality

$$(2.9) \quad \begin{aligned} B_n^{U^2D}(\sigma) &= A_{j-1}^{U^2D}(\text{red}(\sigma_1 \cdots \sigma_{j-1})) + B_{n-j}^{U^2D}(\sigma_{j+1} \cdots \sigma_n), \quad \text{if } \sigma_j = n, \\ &\text{for } j \in [n-1], n \geq 2, \end{aligned}$$

along with the fact noted above that under the conditioned measure $P_n^{\text{av}(132)}|\{\sigma_j = n\}$, the permutation $\text{red}(\sigma_1 \cdots \sigma_{j-1}) \in S_{j-1}$ has the distribution $P_{j-1}^{\text{av}(132)}$, the permutation $\sigma_{j+1} \cdots \sigma_n \in S_{n-j}$ has the distribution $P_{n-j}^{\text{av}(132)}$, and these two permutations are independent. Rather than give a formal proof of (2.9), we convince the reader of its validity by giving an example and then a generic explanation.

Let $\sigma = 435768921$. Then $n = 9$ and $j = 7$. We have $A_{j-1}^{U^2D}(\text{red}(\sigma_1 \cdots \sigma_{j-1})) = A_6^{U^2D}(\text{red}(435768)) = A^{U^2D}(213546) = 2$, because the subsequence 23546 (as well as 13546) corresponds to $UUDU$. We have $B_{n-j}^{U^2D}(\sigma_{j+1} \cdots \sigma_n) = B_2^{U^2D}(21) = 0$. And we have $B_n^{U^2D}(\sigma) = B_9^{U^2D}(435768921) = 2$ because the subsequence 4576892 (as well as several others) corresponds to $UUDUUD$.

Generically, $B_n^{U^2D}(\sigma)$ is the sum of two terms. One of the terms is $A_{j-1}^{U^2D}(\text{red}(\sigma_1 \cdots \sigma_{j-1}))$, which counts the number of full sets of UUD and then adds one for an extra U . This extra U , along with $\sigma_j = n$ and σ_{j+1} supply an additional full set UUD which was counted by $A_{j-1}^{U^2D}(\text{red}(\sigma_1 \cdots \sigma_{j-1}))$ (via the adding one for the extra U). The other term is $B_{n-j}^{U^2D}(\sigma_{j+1} \cdots \sigma_n)$, which counts the remaining sets of UUD .

The second line of (2.7) is obtained using the following rather obvious equality instead of (2.9):

$$B_n^{U^2D}(\sigma) = B_{n-1}^{U^2D}(\sigma_1 \cdots \sigma_{n-1}), \quad \text{if } \sigma_n = n.$$

The second line of (2.8) follows from the equality

$$(2.10) \quad A_n^{U^2D}(\sigma) = \left(A_{j-1}^{U^2D}(\text{red}(\sigma_1 \cdots \sigma_{j-1})) + A_{n-j}^{U^2D}(\sigma_{j+1} \cdots \sigma_n) \right) 1_{A_{n-j}^{U^2D}(\sigma_{j+1} \cdots \sigma_n) \neq 0} + \\ \left(B_{j-1}^{U^2D}(\text{red}(\sigma_1 \cdots \sigma_{j-1})) + 1 \right) 1_{A_{n-j}^{U^2D}(\sigma_{j+1} \cdots \sigma_n) = 0}, \text{ if } \sigma_j = n, \text{ for } j \in [n-1], n \geq 2,$$

along with the fact noted above that under the conditioned measure $P_n^{\text{av}(132)}|\{\sigma_j = n\}$, the permutation $\text{red}(\sigma_1 \cdots \sigma_{j-1}) \in S_{j-1}$ has the distribution $P_{j-1}^{\text{av}(132)}$, the permutation $\sigma_{j+1} \cdots \sigma_n \in S_{n-j}$ has the distribution $P_{n-j}^{\text{av}(132)}$, and these two permutations are independent.

In the case that $A_{n-j}^{U^2D}(\sigma_{j+1} \cdots \sigma_n) \neq 0$, (2.10) is obtained by reasoning similar to that for (2.9). We explain (2.10) in the case that $A_{n-j}^{U^2D}(\sigma_{j+1} \cdots \sigma_n) = 0$ with an example. Let $\sigma = 435786921$ (slightly different than the σ used above). So $n = 9$ and $j = 7$. We have $A_{n-j}^{U^2D}(\sigma_{j+1} \cdots \sigma_n) = A_2^{U^2D}(21) = 0$. We have $B_{j-1}^{U^2D}(\text{red}(\sigma_1 \cdots \sigma_{j-1})) = B_6^{U^2D}(\text{red}(435786)) = B_6^{U^2D}(213564) = 1$ because the subsequence 2354 (as well as several others) corresponds to UUD . And we have $A_n^{U^2D}(\sigma) = A_9^{U^2D}(435786921) = 2$ because the subsequence 45769 (as well as several others) corresponds to $UUDU$. (Note that $A_{j-1}^{U^2D}(\text{red}(\sigma_1 \cdots \sigma_{j-1})) = A_6^{U^2D}(\text{red}(435786)) = A_6^{U^2D}(213564) = 1$ because the subsequence 23 (as well as several others) corresponds to U . Thus, when $A_{n-j}^{U^2D}(\sigma_{j+1} \cdots \sigma_n) = 0$, it is not true in general that $A_n^{U^2D}(\sigma) = A_{j-1}^{U^2D}(\text{red}(\sigma_1 \cdots \sigma_{j-1})) + A_{n-j}^{U^2D}(\sigma_{j+1} \cdots \sigma_n)$.)

The first line of (2.8) is obtained using the following rather obvious equality instead of (2.10):

$$A_n^{U^2D}(\sigma) = A_{n-1}^{U^2D}(\sigma_2 \cdots \sigma_n), \text{ if } \sigma_1 = n.$$

□

From (2.5) and (2.7), it follows that

$$(2.11) \quad b_n = E_n^{\text{av}(132)} B_n^{U^2D} = \sum_{j=1}^n E_n^{\text{av}(132)} (B_n^{U^2D} | \sigma_j = n) P_n^{\text{av}(132)}(\sigma_j = n) = \\ \sum_{j=1}^{n-1} \left(E_{j-1}^{\text{av}(132)} A_{j-1}^{U^2D} + E_{n-j}^{\text{av}(132)} B_{n-j}^{U^2D} \right) \frac{C_{j-1} C_{n-j}}{C_n} + E_{n-1}^{\text{av}(132)} B_{n-1}^{U^2D} \frac{C_{n-1} C_0}{C_n} = \\ \sum_{j=1}^{n-1} (a_{j-1} + b_{n-j}) \frac{C_{j-1} C_{n-j}}{C_n} + b_{n-1} \frac{C_{n-1} C_0}{C_n}, \quad n \geq 2.$$

Multiplying both sides of (2.11) by $C_n t^n$, summing over n from 4 to ∞ , and using (2.6), we obtain

$$(2.12) \quad \begin{aligned} \mathcal{B}^{U^2D}(t) &= \sum_{n=4}^{\infty} C_n b_n t^n = t \sum_{n=4}^{\infty} \left(\sum_{j=1}^{n-1} (a_{j-1} + b_{n-j}) C_{j-1} C_{n-j} \right) t^{n-1} + \\ & t \sum_{n=4}^{\infty} b_{n-1} C_{n-1} t^{n-1}. \end{aligned}$$

Straightforward algebraic calculations along with (2.6) show that

$$(2.13) \quad \begin{aligned} \sum_{n=4}^{\infty} \left(\sum_{j=1}^{n-1} a_{j-1} C_{j-1} C_{n-j} \right) t^{n-1} &= \mathcal{A}^{U^2D}(t) (C(t) - 1); \\ \sum_{n=4}^{\infty} \left(\sum_{j=1}^{n-1} b_{n-j} C_{j-1} C_{n-j} \right) t^{n-1} &= \mathcal{B}^{U^2D}(t) C(t). \end{aligned}$$

From (2.12) and (2.13), we obtain

$$\mathcal{B}^{U^2D}(t) = t \left(\mathcal{A}^{U^2D}(t) (C(t) - 1) + \mathcal{B}^{U^2D}(t) C(t) + \mathcal{B}^{U^2D}(t) \right),$$

which we write as

$$(2.14) \quad \mathcal{B}^{U^2D}(t) = \frac{t (C(t) - 1) \mathcal{A}^{U^2D}(t)}{1 - t - tC(t)}.$$

Note that for $l \in \mathbb{N}$ and $\sigma \in S_l(132)$, $A_l^{U^2D}(\sigma) = 0$ only for $\sigma = l \cdots 21$; thus $P_l^{\text{av}(132)}(A_l^{U^2D} = 0) = \frac{1}{C_l}$. Using this with (2.5) and (2.8), it follows that

$$(2.15) \quad \begin{aligned} a_n &= E_n^{\text{av}(132)} A_n^{U^2D} = \sum_{j=1}^n E_n^{\text{av}(132)} (\mathcal{A}_n^{U^2D} | \sigma_j = n) P_n^{\text{av}(132)} (\sigma_j = n) = \\ & \frac{C_0 C_{n-1}}{C_n} a_{n-1} + \sum_{j=2}^n \left(a_{j-1} \left(1 - \frac{1}{C_{n-j}} \right) + a_{n-j} \right) \frac{C_{j-1} C_{n-j}}{C_n} + \\ & \sum_{j=2}^n \frac{b_{j-1} + 1}{C_{n-j}} \frac{C_{j-1} C_{n-j}}{C_n}. \end{aligned}$$

Multiplying both sides of (2.15) by $C_n t^n$, summing over n from 2 to ∞ and using (2.6), we obtain

$$(2.16) \quad \begin{aligned} \mathcal{A}^{U^2 D}(t) &= \sum_{n=2}^{\infty} C_n a_n t^n = \\ &t \sum_{n=2}^{\infty} C_{n-1} a_{n-1} t^{n-1} + t \sum_{n=2}^{\infty} \left(\sum_{j=2}^n a_{j-1} C_{j-1} C_{n-j} \right) t^{n-1} - t \sum_{n=2}^{\infty} \left(\sum_{j=2}^n a_{j-1} C_{j-1} \right) t^{n-1} + \\ &t \sum_{n=2}^{\infty} \left(\sum_{j=2}^n C_{j-1} a_{n-j} C_{n-j} \right) t^{n-1} + t \sum_{n=2}^{\infty} \left(\sum_{j=2}^n b_{j-1} C_{j-1} \right) t^{n-1} + t \sum_{n=2}^{\infty} \left(\sum_{j=2}^n C_{j-1} \right) t^{n-1}. \end{aligned}$$

Straightforward algebraic calculations along with (2.6) show that

$$(2.17) \quad \begin{aligned} \sum_{n=2}^{\infty} \left(\sum_{j=2}^n a_{j-1} C_{j-1} C_{n-j} \right) t^{n-1} &= \mathcal{A}^{U^2 D}(t) C(t); \\ \sum_{n=2}^{\infty} \left(\sum_{j=2}^n C_{j-1} a_{n-j} C_{n-j} \right) t^{n-1} &= \mathcal{A}^{U^2 D}(t) (C(t) - 1); \\ \sum_{n=2}^{\infty} \left(\sum_{j=2}^n a_{j-1} C_{j-1} \right) t^{n-1} &= \frac{\mathcal{A}^{U^2 D}(t)}{1-t}; \\ \sum_{n=2}^{\infty} \left(\sum_{j=2}^n b_{j-1} C_{j-1} \right) t^{n-1} &= \frac{\mathcal{B}^{U^2 D}(t)}{1-t}; \\ \sum_{n=2}^{\infty} \left(\sum_{j=2}^n C_{j-1} \right) t^{n-1} &= \frac{C(t) - 1}{1-t}. \end{aligned}$$

From (2.16) and (2.17), we obtain

$$(2.18) \quad \begin{aligned} \mathcal{A}^{U^2 D}(t) &= t \left(\mathcal{A}^{U^2 D}(t) + \mathcal{A}^{U^2 D}(t) C(t) - \frac{\mathcal{A}^{U^2 D}(t)}{1-t} + \mathcal{A}^{U^2 D}(t) (C(t) - 1) + \right. \\ &\quad \left. \frac{\mathcal{B}^{U^2 D}(t)}{1-t} + \frac{C(t) - 1}{1-t} \right), \end{aligned}$$

which we write as

$$(2.19) \quad \mathcal{A}^{U^2 D}(t) = \frac{t \left(\mathcal{B}^{U^2 D}(t) + C(t) - 1 \right)}{(1-t)(1-2tC(t)) + t}.$$

Substituting (2.19) in (2.14) and solving for $\mathcal{B}^{U^2D}(t)$, we obtain

$$(2.20) \quad \mathcal{B}^{U^2D}(t) = \frac{t^2 (C(t) - 1)^2}{((1-t)(1-2tC(t)) + t)(1-t-tC(t)) - t^2(C(t) - 1)}.$$

We write the denominator in (2.20) as

$$(2.21) \quad \begin{aligned} & ((1-t)(1-2tC(t)) + t)(1-t-tC(t)) - t^2(C(t) - 1) = \\ & t^2 - t + 1 + (-2t^3 + 3t^2 - 3t)C(t) + 2t^2(1-t)C^2(t). \end{aligned}$$

Using (2.4) and performing some algebra, we have

$$(2.22) \quad \begin{aligned} & t^2 - t + 1 + (-2t^3 + 3t^2 - 3t)C(t) + 2t^2(1-t)C^2(t) = \\ & \frac{1}{2} ((2t^2 - t + 1)\sqrt{1-4t} + (1-t)(1-4t)). \end{aligned}$$

Using (2.4), the numerator in (2.20) can be written as.

$$(2.23) \quad t^2(C(t) - 1)^2 = \frac{1}{2} (2t^2 - 4t + 1 + (2t - 1)\sqrt{1-4t}).$$

From (2.20)-(2.23), we obtain

$$(2.24) \quad \mathcal{B}^{U^2D}(t) = \frac{2t^2 - 4t + 1 + (2t - 1)\sqrt{1-4t}}{(2t^2 - t + 1)\sqrt{1-4t} + (1-t)(1-4t)}.$$

In order to eliminate the square root from the denominator, we multiply the numerator and denominator on the right hand side of (2.24) by the conjugate of the denominator, $-(2t^2 - t + 1)\sqrt{1-4t} + (1-t)(1-4t)$. Performing the algebra, the new denominator can be written as $-4t(1-4t)(t^3 - t + 1)$. Performing the algebra to calculate the new numerator, and dividing the new numerator and the new denominator by $t(1-4t)$, we obtain

$$(2.25) \quad \mathcal{B}^{U^2D}(t) = \frac{N(t)}{D(t)},$$

where

$$(2.26) \quad \begin{aligned} tN(t) &= (2t^2 - 4t + 1)(1-t) + (2t-1)(1-t)(1-4t)^{\frac{1}{2}} \\ &- (2t^2 - 4t + 1)(2t^2 - t + 1)(1-4t)^{-\frac{1}{2}} - (2t-1)(2t^2 - t + 1) \end{aligned}$$

and

$$(2.27) \quad D(t) = -4(t^3 - t + 1).$$

For a function f represented by a power series as $f(t) = \sum_{n=0}^{\infty} f_n t^n$, we use the notation $[t^n]f(t) = f_n$. From [5, p. 381], we have

$$(2.28) \quad [t^n](1-4t)^{-\alpha} = 4^n \frac{n^{\alpha-1}}{\Gamma(\alpha)} \left(1 + O\left(\frac{1}{n}\right) \right), \text{ for } \alpha \in \mathbb{C} - \mathbb{Z}_{\leq 0}.$$

From the transfer theorem [5, Theorem VI.3, p. 390, Example VI.2, p. 395] and (2.28), it follows that if $g(t)$ is analytic in a disk, centered at the origin, of radius larger than $\frac{1}{4}$, then

$$(2.29) \quad [t^n]g(t)(1-4t)^{-\alpha} = g\left(\frac{1}{4}\right)4^n \frac{n^{\alpha-1}}{\Gamma(\alpha)} \left(1 + O\left(\frac{1}{n}\right)\right), \text{ for } \alpha \in \mathbb{C} - \mathbb{Z}_{\leq 0}.$$

From (2.29) it follows immediately that

$$(2.30) \quad [t^n]g(t)t^b(1-4t)^{-\alpha} = g\left(\frac{1}{4}\right)4^{n-b} \frac{n^{\alpha-1}}{\Gamma(\alpha)} \left(1 + O\left(\frac{1}{n}\right)\right), \text{ for } b \in \mathbb{Z} \text{ and } \alpha \in \mathbb{C} - \mathbb{Z}_{\leq 0}.$$

Noting that all of the roots of $D(t)$ have absolute value greater than $\frac{1}{4}$, and applying (2.30) with $\alpha \in \{\frac{1}{2}, -\frac{1}{2}\}$ and $g(t) = -\frac{1}{D(t)}$ to (2.25)-(2.27), it follows that the leading order contribution to $[t^n]\mathcal{B}^{U^2D}(t)$ as $n \rightarrow \infty$ comes from the term $\frac{\frac{1}{4}(2t^2-4t+1)(2t^2-t+1)(1-4t)^{-\frac{1}{2}}}{4(t^3-t+1)} = \frac{4t^3-10t^2+8t-5+t^{-1}}{4(t^3-t+1)}(1-4t)^{-\frac{1}{2}}$. Noting that $\Gamma(\frac{1}{2}) = \sqrt{\pi}$ and $g(\frac{1}{4}) = -\frac{1}{D(\frac{1}{4})} = \frac{16}{49}$ it follows from (2.30) that

$$(2.31) \quad [t^n]\mathcal{B}^{U^2D}(t) \sim \frac{16}{49}4^n \frac{n^{-\frac{1}{2}}}{\sqrt{\pi}} (4 \cdot 4^{-3} - 10 \cdot 4^{-2} + 8 \cdot 4^{-1} - 5 + 4) = \frac{1}{7}4^n \frac{n^{-\frac{1}{2}}}{\sqrt{\pi}}.$$

From (2.3) and (2.2), we have $[t^n]\mathcal{B}^{U^2D}(t) = C_n b_n = C_n E_n^{\text{av}(132)} B_n^{U^2D}$, and as is well known, the Catalan numbers satisfy $C_n \sim 4^n \frac{n^{-\frac{3}{2}}}{\sqrt{\pi}}$. Using these facts with (2.31), we conclude that

$$(2.32) \quad E_n^{\text{av}(132)} B_n^{U^2D} \sim \frac{1}{7}n.$$

Now Theorem 1 follows from (2.32) and (2.1). \square

3. DERIVATION OF THE GENERATING FUNCTIONS FOR THEOREM 2

For $\sigma \in S_n$ and $n \in \mathbb{N}$, define $B_n^{U^3D}(\sigma)$ to be the number of complete sets of $UUUD$ in a longest subsequence in σ of the repeated pattern $UUUD$. Thus,

$$(3.1) \quad B_n^{U^3D}(\sigma) = \begin{cases} \frac{1}{4} \left(L_n^{U^3D}(\sigma) - 1 \right), & \text{if } L_n^{U^3D}(\sigma) \neq 0; \\ 0, & \text{if } L_n^{U^3D}(\sigma) = 0. \end{cases}$$

Also, for convenience, we define $B_0^{U^3D} \equiv 0$.

For $\sigma \in S_n$ and $n \in \mathbb{N}$, define $G_n^{U^3D}(\sigma) = 0$, if $\sigma = n \cdots 21$; otherwise, find a longest subsequence $\{\sigma_{i_j}\}_{j=1}^{4k+2}$, $k \in \mathbb{Z}^+$, of σ for which the up/down pattern is $UUUD \cdots UUUDU$, and define $G_n^{U^3D}(\sigma) = k + 1$. For convenience, we define $G_0^{U^3D} \equiv 0$.

For $\sigma \in S_n$ and $n \in \mathbb{N}$, define $A_n^{U^3D}(\sigma) = 0$, if σ has no increasing subsequence of length three (or equivalently, if σ has no subsequence $\{\sigma_{i_j}\}_{j=1}^3$ which corresponds to the pattern UUU); otherwise, find a longest subsequence $\{\sigma_{i_j}\}_{j=1}^{4k+3}$, $k \in \mathbb{Z}^+$, of σ for which the up/down pattern is $UUUD \cdots UUUDUU$, and define $A_n^{U^3D}(\sigma) = k + 1$. For convenience, we define $A_0^{U^3D} \equiv 0$.

In the sequel, for $j \in \mathbb{N}$, $B_j^{U^3D}$, $G_j^{U^3D}$ and $A_j^{U^3D}$ will always be considered as random variables on the probability space $(S_j(132), P_j^{\text{av}(132)})$. Define

$$(3.2) \quad b_n = E_n^{\text{av}(132)} B_n^{U^3D}; \quad g_n = E_n^{\text{av}(132)} G_n^{U^3D}; \quad a_n = E_n^{\text{av}(132)} A_n^{U^3D},$$

where we have suppressed the notation U^3D .

Define the generating functions for $\{C_n b_n\}_{n=0}^\infty$, $\{C_n g_n\}_{n=0}^\infty$ and $\{C_n a_n\}_{n=0}^\infty$ by

$$(3.3) \quad \begin{aligned} \mathcal{B}^{U^3D}(t) &= \sum_{n=0}^{\infty} C_n b_n t^n; \\ \mathcal{G}^{U^3D}(t) &= \sum_{n=0}^{\infty} C_n g_n t^n; \\ \mathcal{A}^{U^3D}(t) &= \sum_{n=0}^{\infty} C_n a_n t^n. \end{aligned}$$

We will derive a system of three linear equations for $\mathcal{B}^{U^3D}(t)$, $\mathcal{G}^{U^3D}(t)$ and $\mathcal{A}^{U^3D}(t)$ and then solve for one of them explicitly. From the definitions of $B_n^{U^3D}$, $G_n^{U^3D}$ and $A_n^{U^3D}$, we have

$$B_n^{U^3D} \equiv 0, \quad 0 \leq n \leq 4; \quad G_n^{U^3D} \equiv 0, \quad 0 \leq n \leq 1; \quad A_n^{U^3D} = 0, \quad 0 \leq n \leq 2.$$

Thus,

$$(3.4) \quad \begin{aligned} b_n &= 0, \quad 0 \leq n \leq 4; \\ g_n &= 0, \quad 0 \leq n \leq 1; \\ a_n &= 0, \quad 0 \leq n \leq 2. \end{aligned}$$

The following proposition is the key to obtaining a set of three linear equations for the generating functions $\mathcal{B}^{U^3D}(t)$, $\mathcal{G}^{U^3D}(t)$ and $\mathcal{A}^{U^3D}(t)$.

Proposition 2. *i.*

$$(3.5) \quad \begin{aligned} B_n^{U^3D}|\{\sigma_j = n\} &\stackrel{\text{dist}}{=} A_{j-1}^{U^3D} + B_{n-j}^{U^3D}, \quad j \in [n-1], \quad n \geq 2; \\ B_n^{U^3D}|\{\sigma_n = n\} &\stackrel{\text{dist}}{=} B_{n-1}^{U^3D}, \quad n \geq 2, \end{aligned}$$

where on the right hand side of (3.5), $A_{j-1}^{U^3D}$ and $B_{n-j}^{U^3D}$ are understood to be independent.

ii.

(3.6)

$$\begin{aligned} G_n^{U^3D}|\{\sigma_1 = n\} &\stackrel{\text{dist}}{=} G_{n-1}^{U^3D}, \quad n \geq 2; \\ G_n^{U^3D}|\{\sigma_j = n\} &\stackrel{\text{dist}}{=} \left(A_{j-1}^{U^3D} + G_{n-j}^{U^3D}\right) 1_{\{G_{n-j}^{U^3D} \neq 0\}} + \left(B_{j-1}^{U^2D} + 1\right) 1_{\{G_{n-j}^{U^3D} = 0\}}, \\ j &\in \{2, \dots, n\}, \quad n \geq 2, \end{aligned}$$

where on the right hand side of (3.6), $A_{j-1}^{U^3D}$ and $G_{n-j}^{U^3D}$ are understood to be independent and $B_{j-1}^{U^2D}$ and $G_{n-j}^{U^3D}$ are understood to be independent.

iii.

$$\begin{aligned} A_n^{U^3D}|\{\sigma_1 = n\} &\stackrel{\text{dist}}{=} A_{n-1}^{U^3D}, \quad n \geq 2; \\ (3.7) \quad A_n^{U^3D}|\{\sigma_j = n\} &\stackrel{\text{dist}}{=} \left(A_{j-1}^{U^3D} + A_{n-j}^{U^3D}\right) 1_{\{A_{n-j}^{U^3D} \neq 0\}} + G_{j-1}^{U^3D} 1_{\{A_{n-j}^{U^3D} = 0\}}, \\ j &\in \{2, \dots, n\}, \quad n \geq 2, \end{aligned}$$

where on the right hand side of (3.7), $A_{j-1}^{U^3D}$ and $A_{n-j}^{U^3D}$ are understood to be independent and $G_{j-1}^{U^3D}$ and $A_{n-j}^{U^3D}$ are understood to be independent.

Proof. The proof is similar to that of Proposition 1. The first line of (3.5) and the second lines of (3.6) and (3.7) follow from the rather obvious equalities

$$\begin{aligned} B_n^{U^3D}(\sigma) &= B_{n-1}^{U^3D}(\sigma_1 \cdots \sigma_{n-1}), \quad \text{if } \sigma_n = n; \\ G_n^{U^3D}(\sigma) &= G_{n-1}^{U^3D}(\sigma_2 \cdots \sigma_n), \quad \text{if } \sigma_1 = n; \\ A_n^{U^3D}(\sigma) &= A_{n-1}^{U^3D}(\sigma_2 \cdots \sigma_n), \quad \text{if } \sigma_1 = n. \end{aligned}$$

Recall the notation $\text{red}(\sigma)$ that was introduced in the paragraph containing (2.5). The first line of (3.5) follows from the equality

$$\begin{aligned} (3.8) \quad B_n^{U^3D}(\sigma) &= A_{j-1}^{U^3D}(\text{red}(\sigma_1 \cdots \sigma_{j-1})) + B_{n-j}^{U^3D}(\sigma_{j+1} \cdots \sigma_n), \quad \text{if } \sigma_j = n, \\ &\text{for } j \in [n-1], n \geq 2, \end{aligned}$$

along with the fact that under the conditioned measure $P_n^{\text{av}(132)}|\{\sigma_j = n\}$, the permutation $\text{red}(\sigma_1 \cdots \sigma_{j-1}) \in S_{j-1}$ has the distribution $P_{j-1}^{\text{av}(132)}$, the permutation $\sigma_{j+1} \cdots \sigma_n \in S_{n-j}$ has the distribution $P_{n-j}^{\text{av}(132)}$, and these two permutations are independent. The explanation for (3.8) is essentially the same as the explanation for (2.9). Generically, $B_n^{U^3D}(\sigma)$ is the sum of two terms. One of the terms is $A_{j-1}^{U^3D}(\text{red}(\sigma_1 \cdots \sigma_{j-1}))$, which counts the number of full sets of U^3D and then adds one for an extra UU . This extra UU , along

with $\sigma_j = n$ and σ_{j+1} supply an additional full set U^3D which was counted by $A_{j-1}^{U^3D}(\text{red}(\sigma_1 \cdots \sigma_{j-1}))$ (via the adding one for the extra UU). The other term is $B_{n-j}^{U^3D}(\sigma_{j+1} \cdots \sigma_n)$, which counts the remaining sets of U^3D .

The second line of (3.6) follows from the equality

$$(3.9) \quad G_n^{U^3D}(\sigma) = \left(A_{j-1}^{U^3D}(\text{red}(\sigma_1 \cdots \sigma_{j-1})) + G_{n-j}^{U^3D}(\sigma_{j+1} \cdots \sigma_n) \right) 1_{G_{n-j}^{U^3D}(\sigma_{j+1} \cdots \sigma_n) \neq 0} + \left(B_{j-1}^{U^2D}(\text{red}(\sigma_1 \cdots \sigma_{j-1})) + 1 \right) 1_{G_{n-j}^{U^3D}(\sigma_{j+1} \cdots \sigma_n) = 0},$$

if $\sigma_j = n$, for $j \in [n-1], n \geq 2$,

along with the fact that under the conditioned measure $P_n^{\text{av}(132)}|\{\sigma_j = n\}$, the permutation $\text{red}(\sigma_1 \cdots \sigma_{j-1}) \in S_{j-1}$ has the distribution $P_{j-1}^{\text{av}(132)}$, the permutation $\sigma_{j+1} \cdots \sigma_n \in S_{n-j}$ has the distribution $P_{n-j}^{\text{av}(132)}$, and these two permutations are independent. The explanation for (3.9) in the case that $G_{n-j}^{U^3D}(\sigma_{j+1} \cdots \sigma_n) \neq 0$ is similar to the reasoning for (3.8). We explain (3.9) in the case that $G_{n-j}^{U^3D}(\sigma_{j+1} \cdots \sigma_n) = 0$ with an example, the same example used to explain (2.10) in the case that $A_{n-j}^{U^2D}(\sigma_{j+1} \cdots \sigma_n) = 0$. Consider $\sigma = 435786921$. So $n = 9$ and $j = 7$. We have $G_{n-j}^{U^3D}(\sigma_{j+1} \cdots \sigma_n) = G_2^{U^3D}(21) = 0$. We have $B_{j-1}^{U^3D}(\text{red}(\sigma_1 \cdots \sigma_{j-1})) = B_6^{U^3D}(\text{red}(435786)) = B_6^{U^3D}(213564) = 1$ because the subsequence 23564 (as well as 13564) corresponds to U^3D . And we have $G_n^{U^3D}(\sigma) = G_9^{U^3D}(435786921) = 2$ because the subsequence 457869 (as well as 357869) corresponds to U^3DU . (Note that $A_{j-1}^{U^3D}(\text{red}(\sigma_1 \cdots \sigma_{j-1})) = A_6^{U^3D}(\text{red}(435786)) = A_6^{U^3D}(213564) = 1$ because the subsequence 235 (as well as several others) corresponds to UU . Thus, when $G_{n-j}^{U^3D}(\sigma_{j+1} \cdots \sigma_n) = 0$, it is not true in general that $G_n^{U^3D}(\sigma) = A_{j-1}^{U^3D}(\text{red}(\sigma_1 \cdots \sigma_{j-1})) + G_{n-j}^{U^3D}(\sigma_{j+1} \cdots \sigma_n)$.)

The second line of (3.7) follows from the equality

$$(3.10) \quad A_n^{U^3D}(\sigma) = \left(A_{j-1}^{U^3D}(\text{red}(\sigma_1 \cdots \sigma_{j-1})) + A_{n-j}^{U^3D}(\sigma_{j+1} \cdots \sigma_n) \right) 1_{A_{n-j}^{U^3D}(\sigma_{j+1} \cdots \sigma_n) \neq 0} + \left(G_{j-1}^{U^3D}(\text{red}(\sigma_1 \cdots \sigma_{j-1})) \right) 1_{A_{n-j}^{U^3D}(\sigma_{j+1} \cdots \sigma_n) = 0},$$

if $\sigma_j = n$, for $j \in [n-1], n \geq 2$,

along with the fact that under the conditioned measure $P_n^{\text{av}(132)}|\{\sigma_j = n\}$, the permutation $\text{red}(\sigma_1 \cdots \sigma_{j-1}) \in S_{j-1}$ has the distribution $P_{j-1}^{\text{av}(132)}$, the permutation $\sigma_{j+1} \cdots \sigma_n \in S_{n-j}$ has the distribution $P_{n-j}^{\text{av}(132)}$, and these two permutations are independent. The explanation for (3.10) in the case that

$A_{n-j}^{U^3D}(\sigma_{j+1} \cdots \sigma_n) \neq 0$ is similar to the reasoning for (3.8). We explain (3.10) in the case that $A_{n-j}^{U^3D}(\sigma_{j+1} \cdots \sigma_n) = 0$ with an example. Consider $\sigma = 786543921$. So $n = 9$ and $j = 7$. We have $A_{n-j}^{U^3D}(\sigma_{j+1} \cdots \sigma_n) = A_2^{U^3D}(21) = 0$. We have $G_{j-1}^{U^3D}(\text{red}(\sigma_1 \cdots \sigma_{j-1})) = G_6^{U^3D}(\text{red}(786543)) = G_6^{U^3D}(564321) = 1$ because the subsequence 56 corresponds to U . And we have $A_n^{U^3D}(\sigma) = A_9^{U^3D}(786543921) = 1$ because the subsequence 789 corresponds to UU . (Note that $A_{j-1}^{U^3D}(\text{red}(\sigma_1 \cdots \sigma_{j-1})) = A_6^{U^3D}(\text{red}(786543)) = A_6^{U^3D}(564321) = 0$. Thus, the equality $A_n^{U^3D}(\sigma) = A_{j-1}^{U^3D}(\text{red}(\sigma_1 \cdots \sigma_{j-1})) + A_{n-j}^{U^3D}(\sigma_{j+1} \cdots \sigma_n)$ is not true in general when $A_{n-j}^{U^3D}(\sigma_{j+1} \cdots \sigma_n) = 0$.) \square

We now use Proposition 2 to derive a system of three linear equations for $\mathcal{B}^{U^3D}(t)$, $\mathcal{G}^{U^3D}(t)$ and $\mathcal{A}^{U^3D}(t)$. Note from (2.7) and (3.5) that the conditional distributions of $B^{U^2D}(t)$ and $B^{U^3D}(t)$ are exactly the same except that $A^{U^2D}(t)$ in (2.7) is replaced by $A^{U^3D}(t)$ in (3.5). Thus, it follows from (2.14) that

$$(3.11) \quad \mathcal{B}^{U^3D}(t) = \frac{t(C(t) - 1)\mathcal{A}^{U^3D}(t)}{1 - t - tC(t)}.$$

We now turn to $\mathcal{G}^{U^3D}(t)$. Note that for $l \in \mathbb{N}$ and $\sigma \in S_l(132)$, $G_l^{U^3D}(\sigma) = 0$ only for $\sigma = l \cdots 21$; thus $P_l^{\text{av}(132)}(G_l^{U^3D} = 0) = \frac{1}{C_l}$. Using this with (2.5) and (3.6), it follows that

$$(3.12) \quad g_n = E_n^{\text{av}(132)}G_n^{U^3D} = \sum_{j=1}^n E_n^{\text{av}(132)}(G_n^{U^3D} | \sigma_j = n) P_n^{\text{av}(132)}(\sigma_j = n) = \frac{C_0 C_{n-1}}{C_n} g_{n-1} + \sum_{j=2}^n \left(a_{j-1} \left(1 - \frac{1}{C_{n-j}} \right) + g_{n-j} \right) \frac{C_{j-1} C_{n-j}}{C_n} + \sum_{j=2}^n \frac{b_{j-1} + 1}{C_{n-j}} \frac{C_{j-1} C_{n-j}}{C_n}.$$

Multiplying both sides of (3.12) by $C_n t^n$, summing over n from 2 to ∞ and using (3.4), we obtain

$$(3.13) \quad \mathcal{G}^{U^3D}(t) = \sum_{n=2}^{\infty} C_n g_n t^n = t \sum_{n=2}^{\infty} C_{n-1} g_{n-1} t^{n-1} + t \sum_{n=2}^{\infty} \left(\sum_{j=2}^n a_{j-1} C_{j-1} C_{n-j} \right) t^{n-1} - t \sum_{n=2}^{\infty} \left(\sum_{j=2}^n a_{j-1} C_{j-1} \right) t^{n-1} + t \sum_{n=2}^{\infty} \left(\sum_{j=2}^n C_{j-1} g_{n-j} C_{n-j} \right) t^{n-1} + t \sum_{n=2}^{\infty} \left(\sum_{j=2}^n b_{j-1} C_{j-1} \right) t^{n-1} + t \sum_{n=2}^{\infty} \left(\sum_{j=2}^n C_{j-1} \right) t^{n-1}.$$

Considerations almost identical to those that led from (2.16) to (2.18) yield

$$(3.14) \quad \mathcal{G}^{U^3D}(t)(t) = t \left(\mathcal{G}^{U^3D}(t)(t) + \mathcal{A}^{U^3D}(t)(t)C(t) - \frac{\mathcal{A}^{U^3D}(t)(t)}{1-t} + \mathcal{G}^{U^3D}(t)(t)(C(t)-1) + \frac{\mathcal{B}^{U^3D}(t)}{1-t} + \frac{C(t)-1}{1-t} \right),$$

which we write as

$$(3.15) \quad \mathcal{G}^{U^3D}(t)(t) = \frac{\left(tC(t) - \frac{t}{1-t}\right) \mathcal{A}^{U^3D}(t) + \frac{t}{1-t} \mathcal{B}^{U^3D}(t) + \frac{t}{1-t} (C(t)-1)}{1-tC(t)}.$$

We now turn to $\mathcal{A}^{U^3D}(t)$. We need the following lemma.

Lemma 1.

$$(3.16) \quad P_l^{\text{av}(132)}(A_l^{U^3D} = 0) = \frac{2^{l-1}}{C_l}, \quad l \in \mathbb{N}.$$

Proof. For convenience, define

$$(3.17) \quad \begin{aligned} \gamma_l &= P_l^{\text{av}(132)}(A_l^{U^3D} = 0), \quad l \in \mathbb{N}; \\ \gamma_0 &= 1. \end{aligned}$$

For $\sigma \in S_l$, distributed as $P_l^{\text{av}(132)}$, and conditioned on $\sigma_i = l$, the permutations $\text{red}(\sigma_1 \cdots \sigma_{i-1})$ and $\sigma_{i+1} \cdots \sigma_l$ are independent and distributed respectively as $P_{i-1}^{\text{av}(132)}$ and $P_{l-i}^{\text{av}(132)}$. If $\sigma_i = l$, then $A_l^{U^3D}(\sigma) = 0$ if and only if $G_{i-1}^{U^3D}(\text{red}(\sigma_1 \cdots \sigma_{i-1})) = 0$ and $A_{l-i}^{U^3D}(\sigma_{i+1} \cdots \sigma_l) = 0$. Now $G_{i-1}^{U^3D}(\text{red}(\sigma_1 \cdots \sigma_{i-1})) = 0$ if and only if $\text{red}(\sigma_1 \cdots \sigma_{i-1}) = i-1 \cdots 21$. Thus, $P_{i-1}^{\text{av}(132)}(G_{i-1}^{U^3D}(\text{red}(\sigma_1 \cdots \sigma_{i-1})) = 0) = \frac{1}{C_{i-1}}$. Therefore, we have

$$P_l^{\text{av}(132)}(A_l^{U^3D}(\sigma) = 0 | \sigma_i = l) = \frac{\gamma_{l-i}}{C_{i-1}}.$$

Consequently,

$$\gamma_l = P_l^{\text{av}(132)}(A_l^{U^3D} = 0) = \sum_{i=1}^l \frac{C_{i-1} C_{l-i}}{C_l} \frac{\gamma_{l-i}}{C_{i-1}} = \sum_{i=1}^l \frac{C_{l-i} \gamma_{l-i}}{C_l},$$

which we write as

$$(3.18) \quad k_l = \sum_{i=0}^{l-1} k_i, \quad k_i = C_i \gamma_i.$$

Multiply both sides of (3.18) by t^l and write the resulting equation as

$$(3.19) \quad k_l t^l = t \sum_{i=0}^{l-1} k_i t^i t^{l-1-i}.$$

Let $K(t) = \sum_{l=0}^{\infty} k_l t^l$. Summing (3.19) over l from 1 to ∞ , one obtains after some algebra

$$K(t) = 1 + \frac{tK(t)}{1-t},$$

which yields

$$(3.20) \quad K(t) = \frac{1-t}{1-2t} = 1 + \frac{t}{1-2t} = 1 + \sum_{l=1}^{\infty} 2^{l-1} t^l.$$

Thus, $C_l \gamma_l = k_l = 2^{l-1}$, $l \geq 1$. Consequently $P_l^{\text{av}(132)}(A_l^{U^3D} = 0) = \gamma_l = \frac{2^{l-1}}{C_l}$. \square

Using (3.17) with (2.5) and (3.7), it follows that

$$(3.21) \quad \begin{aligned} a_n &= E_n^{\text{av}(132)} A_n^{U^3D} = \sum_{j=1}^n E_n^{\text{av}(132)} (A_n^{U^3D} | \sigma_j = n) P_n^{\text{av}(132)}(\sigma_j = n) = \\ & \frac{C_0 C_{n-1}}{C_n} a_{n-1} + \sum_{j=2}^n (a_{j-1} (1 - \gamma_{n-j}) + a_{n-j}) \frac{C_{j-1} C_{n-j}}{C_n} + \\ & \sum_{j=2}^n g_{j-1} \gamma_{n-j} \frac{C_{j-1} C_{n-j}}{C_n}. \end{aligned}$$

Multiplying both sides of (3.21) by $C_n t^n$ and summing over n from 2 to ∞ , and recalling (3.4), we obtain

$$(3.22) \quad \begin{aligned} \mathcal{A}^{U^3D}(t) &= \sum_{n=2}^{\infty} C_n a_n t^n = t \sum_{n=2}^{\infty} C_{n-1} a_{n-1} t^{n-1} + t \sum_{n=2}^{\infty} \left(\sum_{j=2}^n C_{j-1} a_{j-1} C_{n-j} \right) t^{n-1} - \\ & t \sum_{n=2}^{\infty} \left(\sum_{j=2}^n C_{j-1} a_{j-1} \gamma_{n-j} C_{n-j} \right) t^{n-1} + t \sum_{n=2}^{\infty} \left(\sum_{j=2}^n C_{j-1} C_{n-j} a_{n-j} \right) t^{n-1} + \\ & t \sum_{n=2}^{\infty} \left(\sum_{j=2}^n C_{j-1} g_{j-1} \gamma_{n-j} C_{n-j} \right) t^{n-1}. \end{aligned}$$

By (3.18), $\gamma_{n-j}C_{n-j} = k_{n-j}$. Using this with (3.20), we have

$$\begin{aligned}
 (3.23) \quad & \sum_{n=2}^{\infty} \left(\sum_{j=2}^n C_{j-1} a_{j-1} \gamma_{n-j} C_{n-j} \right) t^{n-1} = \sum_{n=2}^{\infty} \left(\sum_{j=2}^n C_{j-1} a_{j-1} k_{n-j} \right) t^{n-1} = \\
 & K(t) \mathcal{A}^{U^3 D}(t) = \frac{1-t}{1-2t} \mathcal{A}^{U^3 D}(t); \\
 & \sum_{n=2}^{\infty} \left(\sum_{j=2}^n C_{j-1} g_{j-1} \gamma_{n-j} C_{n-j} \right) t^{n-1} = \sum_{n=2}^{\infty} \left(\sum_{j=2}^n C_{j-1} g_{j-1} k_{n-j} \right) t^{n-1} = \\
 & K(t) \mathcal{G}^{U^3 D}(t) = \frac{1-t}{1-2t} \mathcal{G}^{U^3 D}(t).
 \end{aligned}$$

The two terms on the left hand sides of (3.23) appear on the right hand side of (3.22). The other terms on the right hand side of (3.22) can be treated via straightforward algebraic calculations, similar to what was done in previous calculations. This allows for (3.22) to be written term by term as

$$\begin{aligned}
 \mathcal{A}^{U^3 D}(t) &= t \mathcal{A}^{U^3 D}(t) + t \mathcal{A}^{U^3 D}(t) C(t) - \frac{1-t}{1-2t} \mathcal{A}^{U^3 D}(t) + \\
 & t \mathcal{A}^{U^3 D}(t) (C(t) - 1) + \frac{1-t}{1-2t} \mathcal{G}^{U^3 D}(t),
 \end{aligned}$$

which yields

$$(3.24) \quad \mathcal{A}^{U^3 D}(t) = \frac{\frac{t(1-t)}{1-2t} \mathcal{G}^{U^3 D}(t)}{1 - 2tC(t) + \frac{t(1-t)}{1-2t}}.$$

Now (3.11), (3.15) and (3.24) provide a system of three linear equations for the three generating functions $\mathcal{B}^{U^3 D}(t)$, $\mathcal{G}^{U^3 D}(t)$ and $\mathcal{A}^{U^3 D}(t)$. Since $G_n^{U^3 D}(\sigma), A_n^{U^3 D}(\sigma) \in \{B_n^{U^3 D}(\sigma) - 1, B_n^{U^3 D}(\sigma), B_n^{U^3 D}(\sigma) + 1\}$, for all $n \in \mathbb{N}$ and all $\sigma \in S_n$, the leading order asymptotic behavior is the same for $E_n^{\text{av}(132)} B_n^{U^3 D}$, $E_n^{\text{av}(132)} G_n^{U^3 D}$ and $E_n^{\text{av}(132)} A_n^{U^3 D}$. Thus, it doesn't matter which of the generating functions we solve for. We will solve for $\mathcal{G}^{U^3 D}(t)$. We start with (3.15), and replace the term $\mathcal{B}^{U^3 D}(t)$ on the right hand side of (3.15) with the right hand side of (3.11). After rearranging some terms, this gives

$$\begin{aligned}
 (3.25) \quad & \mathcal{G}^{U^3 D}(t) = \frac{t(C(t) - 1)}{(1-t)(1-tC(t))} + \\
 & \left(tC(t) - \frac{t}{1-t} + \frac{t^2(C(t) - 1)}{(1-t)(1-t-tC(t))} \right) \frac{1}{1-tC(t)} \mathcal{A}^{U^3 D}(t).
 \end{aligned}$$

Now we replace $\mathcal{A}^{U^3D}(t)$ on the right hand side of (3.25) with the right hand side of (3.24). This yields an equation in which only the generating function $\mathcal{G}^{U^3D}(t)$ appears. Solving for $\mathcal{G}^{U^3D}(t)$, we obtain

$$(3.26) \quad \mathcal{G}^{U^3D}(t) = \frac{t(C(t) - 1)}{(1 - t)(1 - tC(t))(1 - d_1(t))}, \text{ where}$$

$$d_1(t) = \left(tC(t) - \frac{t}{1 - t} + \frac{t^2(C(t) - 1)}{(1 - t)(1 - t - tC(t))} \right) \left(\frac{1}{1 - tC(t)} \right) \left(\frac{t(1 - t)}{1 - 2t} \right) \times$$

$$\left(\frac{1}{1 - 2tC(t) + \frac{t(1 - t)}{1 - 2t}} \right).$$

4. COMPLETION OF THE PROOF OF THEOREM 2

In (3.26), when we perform the multiplication $(1 - t)(1 - tC(t))d_1(t)$, the second of the four factors in $d_1(t)$ will disappear, and the $1 - t$ in the denominator of two of the terms in the first factor will also disappear. We obtain

$$(4.1) \quad (1 - t)(1 - tC(t))d_1(t) = \left((1 - t)tC(t) - t + \frac{t^2(C(t) - 1)}{1 - t - tC(t)} \right) \left(\frac{t(1 - t)}{1 - 2t} \right) \times$$

$$\left(\frac{1}{1 - 2tC(t) + \frac{t(1 - t)}{1 - 2t}} \right).$$

Multiplying the denominators of the second and third factors on the right hand side of (4.1), we have

$$(4.2) \quad (1 - 2t) \left(1 - 2tC(t) + \frac{t(1 - t)}{1 - 2t} \right) = 1 - t - t^2 - 2tC(t) + 4t^2C(t).$$

Thus, multiplying both the numerator and the denominator on the right hand side of (3.26) by $1 - t - t^2 - 2tC(t) + 4t^2C(t)$, and using (4.1) and (4.2), we obtain

$$(4.3) \quad \mathcal{G}^{U^3D}(t) = \frac{t(C(t) - 1)(1 - t - t^2 - 2tC(t) + 4t^2C(t))}{d_2(t)},$$

where

$$(4.4) \quad d_2(t) = (1 - t)(1 - tC(t))(1 - t - t^2 - 2tC(t) + 4t^2C(t)) -$$

$$\left((1 - t)tC(t) - t + \frac{t^2(C(t) - 1)}{1 - t - tC(t)} \right) t(1 - t).$$

Multiplying the numerator and the denominator on the right hand side of (4.3) by $1 - t - tC(t)$, and using (4.4), we obtain

$$(4.5) \quad \mathcal{G}^{U^3D}(t) = \frac{n(t)}{d(t)},$$

where

$$(4.6) \quad \begin{aligned} n(t) &= t(C(t) - 1) (1 - t - t^2 - 2tC(t) + 4t^2C(t)) (1 - t - tC(t)); \\ d(t) &= (1 - t)(1 - tC(t)) (1 - t - t^2 - 2tC(t) + 4t^2C(t)) (1 - t - tC(t)) - \\ &\quad \left(((1 - t)tC(t) - t) (1 - t - tC(t)) + t^2(C(t) - 1) \right) t(1 - t). \end{aligned}$$

Grouping powers of $C(t)$, we have

$$(4.7) \quad \begin{aligned} n(t) &= A_3(t)C^3(t) + A_2(t)C^2(t) + A_1(t)C(t) + A_0(t); \\ d(t) &= B_3(t)C^3(t) + B_2(t)C^2(t) + B_1(t)C(t) + B_0(t), \end{aligned}$$

where

$$(4.8) \quad \begin{aligned} A_3(t) &= 2t^3 - 4t^2; \quad A_2(t) = t^4 + 5t^3 - 3t^2; \\ A_1(t) &= 4t^4 - 7t^3 + t^2 + t; \quad A_0(t) = -t^4 + 2t^2 - t; \\ B_3(t) &= -4t^5 + 6t^4 - 2t^3; \quad B_2(t) = -2t^5 + 12t^4 - 15t^3 + 5t^2; \\ B_1(t) &= 2t^5 + t^4 - 11t^3 + 12t^2 - 4t; \quad B_0(t) = -t^4 + 3t^2 - 3t + 1. \end{aligned}$$

Recalling the formula for $C(t)$ in (2.4), we have

$$(4.9) \quad \begin{aligned} C^2(t) &= \frac{1 - 2t - \sqrt{1 - 4t}}{2t^2}; \\ C^3(t) &= \frac{1 - 3t - (1 - t)\sqrt{1 - 4t}}{2t^3}. \end{aligned}$$

Letting

$$R := \sqrt{1 - 4t}$$

and substituting from (2.4) and (4.9) in (4.7), we obtain after a lot of algebra

$$(4.10) \quad \begin{aligned} n(t) &= \left(-\frac{(1 - t)A_3(t)}{2t^3} - \frac{A_2(t)}{2t^2} - \frac{A_1(t)}{2t} \right) R + \\ &\quad \frac{(1 - 3t)A_3(t)}{2t^3} + \frac{(1 - 2t)A_2(t)}{2t^2} + \frac{A_1(t)}{2t} + A_0(t); \\ d(t) &= \left(-\frac{(1 - t)B_3(t)}{2t^3} - \frac{B_2(t)}{2t^2} - \frac{B_1(t)}{2t} \right) R + \\ &\quad \frac{(1 - 3t)B_3(t)}{2t^3} + \frac{(1 - 2t)B_2(t)}{2t^2} + \frac{B_1(t)}{2t} + B_0(t). \end{aligned}$$

Using (4.8), one finds that

$$\begin{aligned}
 (4.11) \quad & -\frac{(1-t)A_3(t)}{2t^3} - \frac{A_2(t)}{2t^2} - \frac{A_1(t)}{2t} = t^2(1-2t); \\
 & \frac{(1-3t)A_3(t)}{2t^3} + \frac{(1-2t)A_2(t)}{2t^2} + \frac{A_1(t)}{2t} + A_0(t) = t^3(1-t); \\
 & -\frac{(1-t)B_3(t)}{2t^3} - \frac{B_2(t)}{2t^2} - \frac{B_1(t)}{2t} = -t^4 - \frac{3}{2}t^3 + \frac{9}{2}t^2 - \frac{5}{2}t + \frac{1}{2}; \\
 & \frac{(1-3t)B_3(t)}{2t^3} + \frac{(1-2t)B_2(t)}{2t^2} + \frac{B_1(t)}{2t} + B_0(t) = 2t^4 - \frac{13}{2}t^3 + \frac{15}{2}t^2 - \frac{7}{2}t + \frac{1}{2}.
 \end{aligned}$$

From (4.10) and (4.11), we have

$$\begin{aligned}
 (4.12) \quad & n(t) = t^2(1-2t)R + t^3(1-t); \\
 & d(t) = \left(-t^4 - \frac{3}{2}t^3 + \frac{9}{2}t^2 - \frac{5}{2}t + \frac{1}{2}\right)R + 2t^4 - \frac{13}{2}t^3 + \frac{15}{2}t^2 - \frac{7}{2}t + \frac{1}{2}.
 \end{aligned}$$

Recall that $R = \sqrt{1-4t}$. In order to eliminate the square root in the denominator $d(t)$ in (4.12), we multiply the numerator and denominator by the denominator's conjugate, $-(-t^4 - \frac{3}{2}t^3 + \frac{9}{2}t^2 - \frac{5}{2}t + \frac{1}{2})R + 2t^4 - \frac{13}{2}t^3 + \frac{15}{2}t^2 - \frac{7}{2}t + \frac{1}{2}$. Calling the resulting numerator and denominator by $\bar{n}(t)$ and $\bar{d}(t)$, this yields

$$\begin{aligned}
 (4.13) \quad & \bar{n}(t) = \left(-t^6 - \frac{9}{2}t^5 + 21t^4 - \frac{57}{2}t^3 + \frac{35}{2}t^2 - 5t + \frac{1}{2}\right)\sqrt{1-4t} + \\
 & 6t^6 + \frac{29}{2}t^5 - 58t^4 + \frac{119}{2}t^3 - \frac{55}{2}t^2 + 6t - \frac{1}{2}; \\
 & \bar{d}(t) = 4t^7 + 15t^6 - 56t^5 + 45t^4 + 4t^3 - 19t^2 + 8t - 1.
 \end{aligned}$$

The new denominator factors as

$$\begin{aligned}
 (4.14) \quad & \bar{d}(t) = 4t^7 + 15t^6 - 56t^5 + 45t^4 + 4t^3 - 19t^2 + 8t - 1 = \\
 & (1-4t)(1-t)^2(-t^4 - 6t^3 + 2t^2 + 2t - 1).
 \end{aligned}$$

The two polynomials in the new numerator $\bar{n}(t)$ factor as

$$\begin{aligned}
 (4.15) \quad & -t^6 - \frac{9}{2}t^5 + 21t^4 - \frac{57}{2}t^3 + \frac{35}{2}t^2 - 5t + \frac{1}{2} = \\
 & (1-t)^2\left(-t^4 - \frac{13}{2}t^3 + 9t^2 - 4t + \frac{1}{2}\right); \\
 & 6t^6 + \frac{29}{2}t^5 - 58t^4 + \frac{119}{2}t^3 - \frac{55}{2}t^2 + 6t - \frac{1}{2} = \\
 & (1-4t)(1-t)\left(\frac{3}{2}t^4 + \frac{11}{2}t^3 - 8t^2 + \frac{7}{2}t - \frac{1}{2}\right).
 \end{aligned}$$

From (4.13)-(4.15) and (4.5), we conclude that

$$(4.16) \quad \mathcal{G}^{U^3D}(t) = \frac{-t^4 - \frac{13}{2}t^3 + 9t^2 - 4t + \frac{1}{2}}{-t^4 - 6t^3 + 2t^2 + 2t - 1} (1 - 4t)^{-\frac{1}{2}} + \frac{\frac{3}{2}t^4 + \frac{11}{2}t^3 - 8t^2 + \frac{7}{2}t - \frac{1}{2}}{(1-t)(-t^4 - 6t^3 + 2t^2 + 2t - 1)}.$$

The smallest absolute value among the roots of $-t^4 - 6t^3 + 2t^2 + 2t - 1$ is larger than $\frac{1}{4}$; thus, $\frac{1}{-t^4 - 6t^3 + 2t^2 + 2t - 1}$ and $\frac{1}{(1-t)(-t^4 - 6t^3 + 2t^2 + 2t - 1)}$ are analytic in a ball centered at the origin of radius larger than $\frac{1}{4}$. Thus, applying (2.30) in the case $\alpha = \frac{1}{2}$ with $g(t) = \frac{1}{-t^4 - 6t^3 + 2t^2 + 2t - 1}$ and with $g(t) = \frac{1}{(1-t)(-t^4 - 6t^3 + 2t^2 + 2t - 1)}$, it follows from (4.16) that the leading order asymptotic contribution to $[t^n]\mathcal{G}^{U^3D}(t)$ comes from the term $\frac{-t^4 - \frac{13}{2}t^3 + 9t^2 - 4t + \frac{1}{2}}{-t^4 - 6t^3 + 2t^2 + 2t - 1} (1 - 4t)^{-\frac{1}{2}}$. Since $\frac{1}{-t^4 - 6t^3 + 2t^2 + 2t - 1}|_{t=\frac{1}{4}} = -\frac{256}{121}$, we conclude from (2.30) that

$$(4.17) \quad [t^n]\mathcal{G}^{U^3D}(t) \sim \frac{256}{121} 4^n \frac{n^{-\frac{1}{2}}}{\sqrt{\pi}} \left(4^{-4} + \frac{13}{2} \cdot 4^{-3} - 9 \cdot 4^{-2} + 4 \cdot 4^{-1} - \frac{1}{2} \right) = \frac{256}{121} 4^n \frac{n^{-\frac{1}{2}}}{\sqrt{\pi}} \frac{11}{256} = \frac{1}{11} 4^n \frac{n^{-\frac{1}{2}}}{\sqrt{\pi}}.$$

From (3.3) and (3.2), we have $[t^n]\mathcal{G}^{U^3D}(t) = C_n g_n = C_n E_n^{\text{av}(132)} G_n^{U^3D}$. As previously noted, the Catalan numbers satisfy $C_n \sim 4^n \frac{n^{-\frac{3}{2}}}{\sqrt{\pi}}$. Using these facts with (4.17), we conclude that

$$(4.18) \quad E_n^{\text{av}(132)} G_n^{U^3D} \sim \frac{1}{11} n.$$

Theorem 2 now follows from (4.18), (3.1) and the fact that the leading order asymptotic behavior of $E_n^{\text{av}(132)} B_n^{U^3D}$ and of $E_n^{\text{av}(132)} G_n^{U^3D}$ coincide. \square

REFERENCES

- [1] Abrams, A., Babson, E., Landau, H., Landau, Z. and Pommersheim, J., *A Central Limit Theorem for Repeating Patterns*, preprint (2024).
- [2] Baik, J., Deift, P. and Johansson, K., *On the distribution of the length of the longest increasing subsequence of random permutations*, J. Amer. Math. Soc. **12** (1999), 1119-1178.
- [3] Bóna, M., *Combinatorics of Permutations*, Discrete Math. Appl. (Boca Raton) CRC Press, Boca Raton, FL, (2012).
- [4] Deutsch, E., Hildebrand, A. J. and Wilf, H. S., *Longest increasing subsequences in pattern-restricted permutations* Electron. J. Combin. **9** (2002), Research paper 12, 8 pp.
- [5] Flajolet, P. and Sedgewick, R., *Analytic Combinatorics*, Cambridge University Press, Cambridge, (2009).

- [6] Firro, G., Mansour, T. and Wilson, M. C., *Longest alternating subsequences in pattern-restricted permutations*, Electron. J. Combin. **14** (2007), Research Paper 34, 17 pp.
- [7] Logan, B. F. and Shepp, L. A., *A variational problem for random Young tableaux* Advances in Math. **26** (1977), 206-222.
- [8] Pinsky, R. G., *Large deviations for the longest alternating and the longest increasing subsequence in a random permutation avoiding a pattern of length three*, Electron. Commun. Probab. **29** (2024), Paper No. 10, 11 pp.
- [9] Romik, D., *The Surprising Mathematics of Longest Increasing Subsequences*, IMS Textb., **4** Cambridge University Press, New York, (2015).
- [10] Simion, R. and Schmidt, F., *Restricted permutations*, European J. Combin. **6** (1985), 383-406.
- [11] Stanley, R., *Longest alternating subsequences of permutations*, Michigan Math. J. **57** (2008), 675-687.
- [12] Veršik, A. M. and Kerov, S. V., *Asymptotic behavior of the Plancherel measure of the symmetric group and the limit form of Young tableaux*, Dokl. Akad. Nauk SSSR **233** (1977), 1024-1027.
- [13] , Widom, H., *On the limiting distribution for the longest alternating sequence in a random permutation*, Electron. J. Combin. **13** (2006), Article R25.

DEPARTMENT OF MATHEMATICS, TECHNION—ISRAEL INSTITUTE OF TECHNOLOGY,
HAIFA, 32000, ISRAEL

Email address: `pinsky@technion.ac.il`

URL: `https://pinsky.net.technion.ac.il/`